



The Laplacian spectral radius of trees and maximum vertex degree

Xi-Ying Yuan^{a,*}, Yue Liu^b, Miaomiao Han^a

^a Department of Mathematics, Shanghai University, Shanghai, 200444, China

^b College of mathematics and computer science, Fuzhou University, Fuzhou, 350108, China

ARTICLE INFO

Article history:

Received 18 June 2010

Received in revised form 1 January 2011

Accepted 16 January 2011

Available online 16 February 2011

Keywords:

Tree

Maximum degree

Laplacian spectral radius

ABSTRACT

Let $\Delta(T)$ and $\mu(T)$ denote the maximum degree and the Laplacian spectral radius of a tree T , respectively. In this paper we prove that for two trees T_1 and T_2 on n ($n \geq 21$) vertices, if $\Delta(T_1) > \Delta(T_2)$ and $\Delta(T_1) \geq \lceil \frac{11n}{30} \rceil + 1$, then $\mu(T_1) > \mu(T_2)$, and the bound “ $\Delta(T_1) \geq \lceil \frac{11n}{30} \rceil + 1$ ” is the best possible. We also prove that for two trees T_1 and T_2 on $2k$ ($k \geq 4$) vertices with perfect matchings, if $\Delta(T_1) > \Delta(T_2)$ and $\Delta(T_1) \geq \lceil \frac{k}{2} \rceil + 2$, then $\mu(T_1) > \mu(T_2)$.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Let $G = (V(G), E(G))$ be a simple graph. Denote by $N(v)$ the set of all the neighbors of a vertex v in G , and by $d(v)$ the degree of v . Denote by $m(v)$ the average of the degrees of the vertices adjacent to v , i.e., $m(v) = \frac{\sum_{u \in N(v)} d(u)}{d(v)}$.

Let $D(G)$ be the diagonal matrix of vertex degrees. The Laplacian matrix $L(G)$ of G is defined by $L(G) = D(G) - A(G)$, where $A(G)$ is the adjacency matrix of G . In this paper, the characteristic polynomial $\det(xI - L(G))$ is denoted by $\Phi(L(G); x)$ (or simply $\Phi(L(G))$). We always use $\mu_i(B)$ to denote the i th eigenvalue of a real symmetric matrix B (of order n), and the inequalities $\mu_1(B) \geq \mu_2(B) \geq \dots \geq \mu_n(B)$ hold. Write $\mu_i(G) = \mu_i(L(G))$ for short. The eigenvalue $\mu_1(G)$ is called the Laplacian spectral radius of G , denoted by $\mu(G)$. We may cite some recent papers (see [5,6,12,13,20]) on the ordering of trees based on the Laplacian spectral radius and Laplacian coefficients.

Denote by $\Delta(G)$ for the maximum degree of G . Let $\mathcal{T}(n)$ be the set of trees on n vertices. It is proved in [18] that the Laplacian spectral radius of a tree T in $\mathcal{T}(n)$ ($n \geq 6$) strictly increases with its maximum degree when $\Delta(T) \geq \lceil \frac{n}{2} \rceil + 1$. In this paper, we prove that the bound “ $\Delta(T) \geq \lceil \frac{n}{2} \rceil + 1$ ” can be improved as “ $\Delta(T) \geq \lceil \frac{11n}{30} \rceil + 1$ ”, and we point out that the bound “ $\Delta(T) \geq \lceil \frac{11n}{30} \rceil + 1$ ” is the best possible.

Let $\mathbf{T}(2k)$ be the set of trees on $2k$ vertices with perfect matchings, and $\mathbf{T}(2k, \Delta) = \{T \in \mathbf{T}(2k) \mid \Delta(T) = \Delta\}$. In this paper, we determine the unique tree that takes the largest values of $\mu(T)$ of the trees T in $\mathbf{T}(2k, \Delta)$, when $\Delta \geq \lceil \frac{k}{2} \rceil + 1$ (see Theorem 3.1). For two trees T_1 and T_2 in $\mathbf{T}(2k)$ ($k \geq 4$) we prove that if $\Delta(T_1) > \Delta(T_2)$ and $\Delta(T_1) \geq \lceil \frac{k}{2} \rceil + 2$, then $\mu(T_1) > \mu(T_2)$.

2. A relation between $\mu(T)$ and $\Delta(T)$ of trees in $\mathcal{T}(n)$

Let $\mathcal{T}(n)$ be the set of trees on n vertices, and $\mathcal{T}(n, \Delta) = \{T \in \mathcal{T}(n) \mid \Delta(T) = \Delta\}$. Denote by T^* the tree with the largest Laplacian spectral radius among all the trees in $\mathcal{T}(n, \Delta)$. The following characterization for T^* is due to Yu (see [19]).

* Corresponding author.

E-mail address: xiyingyuan2007@hotmail.com (X.-Y. Yuan).




Fig. 1. The tree $F(n, \Delta)$.

Lemma 2.1 ([19]). Let T^* be a tree with the largest Laplacian spectral radius among all the trees in $\mathcal{T}(n, \Delta)$. Then

- (1) it is a rooted tree with c as a root;
- (2) its height with respect to c is equal to h (for some h);
- (3) each pendant vertex of T^* is at distance $h - 1$ or h from c ;
- (4) each vertex, except possible one, is of degree 1 or Δ ;
- (5) the vertex of degree d ($1 < d < \Delta$), if any, is at distance $h - 1$ from c .

It is easy to see that the vertex c of T^* has maximum degree. The height $h = 1$ if and only if $\Delta = n - 1$, and $h = 2$ if and only if $\sqrt{n-1} \leq \Delta \leq n - 2$. When the height of T^* is at most 2, then T^* is determined by Lemma 2.1 up to isomorphism. When $\lceil \frac{n+6}{3} \rceil \leq \Delta \leq \lceil \frac{n}{2} \rceil - 1$ ($n \geq 18$) the tree T^* has height 2, so we may obtain the following result by Lemma 2.1.

Corollary 2.1. Let T be a tree in $\mathcal{T}(n, \Delta)$, and $F(n, \Delta)$ be the tree as shown in Fig. 1. When $\lceil \frac{n+1}{3} \rceil \leq \Delta \leq \lceil \frac{n}{2} \rceil - 1$ ($n \geq 18$), we have $\mu(T) \leq \mu(F(n, \Delta))$, and equality holds if and only if $T \cong F(n, \Delta)$.

The tree $S(n, \Delta)$ on n vertices is called a double star graph, which is obtained by joining the center of $K_{1, \Delta-1}$ to that of $K_{1, n-\Delta-1}$ by an edge, where $\Delta \geq \lceil \frac{n}{2} \rceil$. Let $G_1 u : v G_2$ be the graph obtained by joining the vertex u of the graph G_1 to the vertex v of the graph G_2 by an edge, where G_1 and G_2 are disjoint. Then we have $F(n, \Delta) \cong S(2\Delta - 1, \Delta) v : w K_{1, n-2\Delta}$, where $F(n, \Delta)$, v and w are shown in Fig. 1.

If $v \in V(G)$, let $L_v(G)$ be the principal sub-matrix of $L(G)$ obtained by deleting the row and column corresponding to the vertex v . The following result is due to Guo [9].

Lemma 2.2 ([9]). Let $G \cong G_1 u : v G_2$. Then

$$\Phi(L(G)) = \Phi(L(G_1))\Phi(L(G_2)) - \Phi(L(G_1))\Phi(L_v(G_2)) - \Phi(L(G_2))\Phi(L_u(G_1)).$$

By using Lemma 2.2 recursively, we may give the characteristic polynomial of matrix $L(F(n, \Delta))$. Let

$$\begin{aligned} f(x) = & x^5 - (n+4)x^4 + (2\Delta n + 2n - 3\Delta^2 + 2\Delta + 7)x^3 - (\Delta^2 n + 2\Delta n + 3n - 2\Delta^3 - 2\Delta^2 + 2\Delta + 6)x^2 \\ & + (3\Delta n + n - 5\Delta^2 + 4\Delta + 2)x - n. \end{aligned} \quad (2.1)$$

Lemma 2.3. Let $F(n, \Delta)$ be the graph as shown in Fig. 1, and the function $f(x)$ be defined in (2.1). We have $\Phi(L(F(n, \Delta)); x) = x(x-1)^{n-6}f(x)$.

Proof. First we give the characteristic polynomial of matrix $L(S(2\Delta - 1, \Delta))$ by using Lemma 2.2 (we may see [7] or [16] for the characteristic polynomial of matrix $L(S(2\Delta - 1, \Delta))$). As we all know that $\Phi(L(K_{1, \Delta}); x) = x(x - \Delta - 1)(x - 1)^{\Delta-1}$, and $\Phi(L_u(K_{1, \Delta}); x) = (x - 1)^\Delta$, where u is the center of the star $K_{1, \Delta}$. Now take the non-pendant edge of $S(2\Delta - 1, \Delta)$ as the edge uv of G in Lemma 2.2. We have

$$\begin{aligned} \Phi(L(S(2\Delta - 1, \Delta)); x) &= x^2(x - \Delta)(x - \Delta + 1)(x - 1)^{2\Delta-5} - x(x - \Delta)(x - 1)^{2\Delta-4} - x(x - \Delta + 1)(x - 1)^{2\Delta-4} \\ &= x(x - 1)^{2\Delta-5}[x^3 - (2\Delta + 1)x^2 + (\Delta^2 + \Delta + 1)x - (2\Delta - 1)]. \end{aligned}$$

It is easy to check that $\Phi(L_v(S(2\Delta - 1, \Delta)); x) = (x - 1)^{2\Delta-4}[x^2 - (\Delta + 1)x + 1]$, where v is the vertex of the double star graph $S(2\Delta - 1, \Delta)$ with degree $\Delta - 1$.

Now we give the characteristic polynomial of matrix $L(F(n, \Delta))$. Take the edge vw of $F(n, \Delta)$ as the edge uv of G in Lemma 2.2.

$$\begin{aligned} \Phi(L(F(n, \Delta)); x) &= x^2(x - 1)^{n-6}(x - n + 2\Delta - 1)[x^3 - (2\Delta + 1)x^2 + (\Delta^2 + \Delta + 1)x - (2\Delta - 1)] \\ &\quad - x(x - 1)^{n-5}[x^3 - (2\Delta + 1)x^2 + (\Delta^2 + \Delta + 1)x - (2\Delta - 1)] \\ &\quad - x(x - 1)^{n-5}(x - n + 2\Delta - 1)[x^2 - (\Delta + 1)x + 1] \\ &= x(x - 1)^{n-6}f(x). \quad \square \end{aligned}$$

Lemma 2.4 ([7]). Let G be a connected graph on n vertices having at least one edge, then $\mu(G) \geq \Delta(G) + 1$, with equality if and only if $\Delta(G) = n - 1$.

The following inequalities are known as Cauchy's inequalities.

Lemma 2.5. Let H be a Hermitian matrix of order n and H' be a principal submatrix of H of order m . Then the inequalities $\mu_{n-m+i}(H) \leq \mu_i(H') \leq \mu_i(H)$ ($i = 1, 2, \dots, m$) hold.

Lemma 2.6. When $\lceil \frac{11n}{30} \rceil \leq \Delta \leq \lceil \frac{n}{2} \rceil - 1$ ($n \geq 10$), we have $\mu(F(n, \Delta)) < \Delta + 2$.

Proof. First we prove that $\mu_2(F(n, \Delta)) \leq \Delta + 1$. Using Lemma 2.5 twice, we have

$$\begin{aligned} \mu_2(F(n, \Delta)) &\leq \mu_1(L_v(F(n, \Delta))) \\ &\leq \max\{\mu_1(K_{1,\Delta}), \mu_1(K_{1,n-2\Delta+1})\} \\ &= \Delta + 1, \end{aligned}$$

where v is the vertex of $F(n, \Delta)$ as shown in Fig. 1. So $\mu_2(F(n, \Delta)) \leq \Delta + 1$ holds.

From Lemma 2.3 we have

$$\Phi(L(F(n, \Delta)); x) = x(x-1)^{n-6}f(x). \quad (2.2)$$

From (2.2) we know that $\mu(F(n, \Delta))$ is the largest root of the equation $f(x) = 0$. And when $\lceil \frac{11n}{30} \rceil \leq \Delta \leq \lceil \frac{n}{2} \rceil - 1$, it is easy to verify that

$$f(\Delta + 1) = -[\Delta^2(3\Delta - n - 1) + 2(n - 2\Delta)] < 0, \quad (2.3)$$

$$f(\Delta + 2) = 4 + (30\Delta - 11n) + (14\Delta^2 - 5\Delta n) > 0. \quad (2.4)$$

From (2.3) and (2.4) we know that there exists a root of the equation $f(x) = 0$ in the interval $(\Delta + 1, \Delta + 2)$. Note that $\mu_2(F(n, \Delta)) \leq \Delta + 1$, so we have $\mu(F(n, \Delta)) < \Delta + 2$. \square

Lemma 2.7 ([14]). Let G be a graph on n vertices. Then $\mu(G) \leq \max\{d(v) + m(v) \mid v \in V(G)\}$.

Lemma 2.8. Let T be a tree on n ($n \geq 21$) vertices with $\Delta(T) \leq \lceil \frac{11n}{30} \rceil - 1$. Then $\mu(T) < \lceil \frac{11n}{30} \rceil + 2$.

Proof. Let v be a vertex of tree T and write $d(v) = s$. Now we distinguish the following two cases depending on the value of s .

Case 1. $s \leq 2$.

$$d(v) + m(v) \leq 2 + \Delta(T) \leq 2 + \left\lceil \frac{11n}{30} \right\rceil - 1 < \left\lceil \frac{11n}{30} \right\rceil + 2. \quad (2.5)$$

Case 2. $3 \leq s \leq \lceil \frac{11n}{30} \rceil - 1$.

Denote by $|E(T)|$ the number of edges of the tree T . Since tree T is a bipartite graph and all neighbors of v belong to exactly one partition, we have

$$m(v) = \frac{1}{s} \sum_{u \in N(v)} d(u) \leq \frac{1}{s} |E(T)| = \frac{n-1}{s}.$$

Hence

$$d(v) + m(v) \leq s + \frac{n-1}{s}.$$

Let $g(s) = s + \frac{n-1}{s}$. Then $g''(s) = \frac{2(n-1)}{s^3} > 0$, and so $g(s)$ is convex when $s > 0$ and $n \geq 2$. Hence when $3 \leq s \leq \lceil \frac{11n}{30} \rceil - 1$, we have

$$g(s) \leq \max \left\{ g(3), g \left(\left\lceil \frac{11n}{30} \right\rceil - 1 \right) \right\} = \max \left\{ 3 + \frac{n-1}{3}, \left\lceil \frac{11n}{30} \right\rceil - 1 + \frac{n-1}{\left\lceil \frac{11n}{30} \right\rceil - 1} \right\}.$$

When $n \geq 21$, it is easy to check that $3 + \frac{n-1}{3} < \lceil \frac{11n}{30} \rceil + 2$, and $\frac{n-1}{\lceil \frac{11n}{30} \rceil - 1} < 3$. So $g(s) < \lceil \frac{11n}{30} \rceil + 2$.

Finally, we have

$$d(v) + m(v) < \left\lceil \frac{11n}{30} \right\rceil + 2. \quad (2.6)$$

Combining (2.5) and (2.6) and Lemma 2.7 we have $\mu(T) < \lceil \frac{11n}{30} \rceil + 2$. \square

The following result was proved in [18] (see Theorem 2.1 and Lemma 3.1 of [18]).

Lemma 2.9 ([18]). Let T be a tree in $\mathcal{T}(n, \Delta)$. When $\Delta \geq \lceil \frac{n}{2} \rceil$ ($n \geq 6$), we have $\mu(T) \leq \mu(S(n, \Delta)) < \Delta + 2$.

Now we are ready to give the main result of this section.

Theorem 2.1. Let $T_1, T_2 \in \mathcal{T}(n)$ ($n \geq 21$). If $\Delta(T_1) > \Delta(T_2)$ and $\Delta(T_1) \geq \lceil \frac{11n}{30} \rceil + 1$, then $\mu(T_1) > \mu(T_2)$.

Proof. (1) If $\Delta(T_2) \geq \lceil \frac{n}{2} \rceil$, from Lemmas 2.4 and 2.9, we have

$$\mu(T_1) \geq \Delta(T_1) + 1 \geq \Delta(T_2) + 2 > \mu(S(n, \Delta(T_2))) \geq \mu(T_2).$$

(2) If $\lceil \frac{11n}{30} \rceil \leq \Delta(T_2) \leq \lceil \frac{n}{2} \rceil - 1$, from Lemma 2.4, Corollary 2.1, and Lemma 2.6 we have

$$\mu(T_1) \geq \Delta(T_1) + 1 \geq \Delta(T_2) + 2 > \mu(F(n, \Delta(T_2))) \geq \mu(T_2).$$

(3) If $\Delta(T_2) \leq \lceil \frac{11n}{30} \rceil - 1$, from Lemmas 2.4 and 2.8 we have

$$\mu(T_1) \geq \Delta(T_1) + 1 \geq \left\lceil \frac{11n}{30} \right\rceil + 2 > \mu(T_2).$$

The proof is completed. \square

Denote by $B(n, \Delta)$ ($2 \leq \Delta \leq n - 1$) the tree on n vertices, which is obtained by joining one pendant vertex of $K_{1,\Delta}$ to one pendant vertex of the path $P_{n-\Delta-1}$ by an edge. We may point out that the tree $B(n, \Delta)$ has simultaneously maximal all Laplacian coefficients among trees with n vertices and maximal vertex degree Δ (see [12]).

Remark 2.1. Theorem 2.1 indicates that the Laplacian spectral radius of a tree T in $\mathcal{T}(n)$ ($n \geq 21$) strictly increases with its maximum degree when $\Delta(T) \geq \lceil \frac{11n}{30} \rceil + 1$. The bound “ $\Delta(T) \geq \lceil \frac{11n}{30} \rceil + 1$ ” is the best possible. To see this, let $T_1 = B(90, 33)$ and $T_2 = F(90, 32)$ (tree $F(n, \Delta)$ is shown in Fig. 1). By using the software *Mathematica* we have $\mu(T_1) = 34.0009$ (up to four decimal places) and $\mu(T_2) = 34.0066$. We have $\Delta(T_1) > \Delta(T_2)$ while $\mu(T_1) < \mu(T_2)$.

3. The tree with the largest Laplacian spectral radius in $\mathbf{T}(2k, \Delta)$

Let $\mathbf{T}(2k, \Delta) = \{T \in \mathbf{T}(2k) \mid \Delta(T) = \Delta\}$. In this section we assume that $\Delta \geq \lceil \frac{k}{2} \rceil + 1$ ($k \geq 4$), then each tree in $\mathbf{T}(2k, \Delta)$ contains exactly one vertex with degree Δ . Let T^* be a tree from $\mathbf{T}(2k, \Delta)$ whose Laplacian spectral radius attains the maximum value. In this section we will characterize the tree T^* . First we formulate some basic tools.

Let $P = v_0 v_1 \cdots v_k$ ($k \geq 1$) be a path of a tree T . If $d_T(v_0) \geq 3$, $d_T(v_k) \geq 3$ and $d_T(v_i) = 2$ ($0 < i < k$), then P is called an *internal path* of T (see [10]).

Lemma 3.1 ([17]). Let uv be an edge on an internal path of tree T . Denote by T^\sharp the tree obtained from T by contracting the edge uv , i.e., identifying the vertices u and v in T . Then $\mu(T^\sharp) > \mu(T)$.

Lemma 3.2 ([8]). Let G be a connected bipartite graph, and G' be a subgraph of G . Then $\mu(G) \geq \mu(G')$, and equality holds if and only if $G \cong G'$.

Lemma 3.3 ([15]). $L(G) = D(G) - A(G)$ and $Q(G) = D(G) + A(G)$ have the same spectrum if and only if G is a bipartite graph.

The matrix $Q(G) = D(G) + A(G)$ is called the signless Laplacian of graph G . See [2–4] for some recent results about the theory of signless Laplacians of graphs. By Lemma 3.3, for any tree T , its Laplacian spectral radius $\mu(T)$ is equal to the largest eigenvalue of $Q(T)$ ($\mu_1(Q(T))$). Since $Q(T)$ is a nonnegative irreducible positive semidefinite symmetric matrix, there is a positive unit eigenvector X belonging to $\mu(T)$. In the following proof, if v is a vertex of a graph G , then x_v denotes the element of X corresponding to v .

Lemma 3.4 ([11]). Let u, v be two vertices of a connected graph G . Suppose v_1, v_2, \dots, v_s ($1 \leq s \leq d(v)$) are some vertices of $N_G(v) \setminus (N_G(u) \cup \{u\})$. Let G' be the graph obtained from G by deleting the edges uv_1, uv_2, \dots, uv_s and adding the edges uv_1, uv_2, \dots, uv_s . Let X be the unit eigenvector of $Q(G)$ corresponding to $\mu_1(Q(G))$. If $x_u \geq x_v$, then $\mu_1(Q(G')) > \mu_1(Q(G))$.

Lemma 3.5 ([19]). Let st and uv be two edges of a tree T , and $sv, tu \notin E(T)$. Let T' be the tree obtained from T by deleting the edges st and uv , and adding edges sv and tu . Let X be the eigenvector of $Q(T)$ corresponding to $\mu(T)$. Then the following holds: If $(x_s - x_u)(x_v - x_t) \geq 0$, then $\mu(T') \geq \mu(T)$, and the equality holds if and only if $x_s = x_u$ and $x_v = x_t$.

In the following proofs, we always suppose that w is the vertex of T^* with degree Δ , and X is the unit eigenvector of $Q(T^*)$ corresponding to $\mu(T^*)$. If a tree T has a perfect matching, then the matching is unique, denoted by $M(T)$ in this paper.

Lemma 3.6. There exists at most one vertex, say u , of T^* with degree $2 < d(u) < \Delta$.

Proof. Suppose to the contrary that there exist two vertices, say u, v , such that $2 < d(u), d(v) < \Delta$. Without loss of generality suppose that $x_u \geq x_v$. The assumption $d(v) > 2$ ensures the existence of a vertex in $N(v)$, say t , such that t does not lie on the (unique) path between v and u , and $tv \notin M(T^*)$. Let

$$T' = T^* - vt + ut,$$

then T' is also in $\mathbf{T}(2k, \Delta)$. Thus $\mu(T') > \mu(T^*)$ follows from Lemma 3.4. This contradicts to the definition of T^* . \square

Lemma 3.7. *If T^* has a vertex, say u , with degree $2 < d(u) < \Delta$, then $uw \in E(T^*)$.*

Proof. Denote by $uu_1 \cdots u_s (=w)$ the (unique) path between u and w . If $u_1 = w$, we are done. So $u_1 \neq w$, and the edge uu_1 lies on some internal path of T^* . We distinguish the following two cases.

Case 1. $uu_1 \in M(T^*)$.

We may contract the edge uu_1 , and then attach a pendant edge to the vertex u , the resulting tree is denoted by T' . It is easy to see that $T' \in \mathbf{T}(2k, \Delta)$. Then $\mu(T') > \mu(T^*)$ follows Lemmas 3.1 and 3.2. This contradicts to the definition of T^* .

Case 2. $uu_1 \notin M(T^*)$.

Suppose $uu' \in M(T^*)$, then $u' \neq w$. Contract the edge uu_1 , and then attach a pendant edge to the vertex u' , the resulting tree is denoted by T'' . Then $T'' \in \mathbf{T}(2k, \Delta)$. So we have $\mu(T'') > \mu(T^*)$, which follows Lemmas 3.1 and 3.2. This contradicts to the definition of T^* .

The proof is completed. \square

Lemma 3.8. *If T^* has a vertex, say u , with degree $2 < d(u) < \Delta$, then $uw \notin M(T^*)$.*

Proof. Suppose to the contrary that $uw \in M(T^*)$. Let w_1 be a neighbor of w different from u , and w_1 is saturated by w_2 ($w_2 \neq w$). And let u_1 be a neighbor of u different from w , and u_1 is saturated by u_2 ($u_2 \neq u$). Then we may apply the following transformation to T^* .

Set

$$T' = \begin{cases} T^* - w_1w_2 + uw_2, & \text{if } x_u \geq x_{w_1}; \\ T^* - u_1u_2 + w_2u_2, & \text{if } x_{w_2} \geq x_{u_1}; \\ T^* - \{w_1w_2, uu_1\} + \{w_1u_1, uw_2\}, & \text{if } x_{w_1} > x_u \text{ and } x_{u_1} > x_{w_2}. \end{cases}$$

Then $\mu(T') > \mu(T^*)$ follows from Lemmas 3.4 and 3.5, while T' is also in $\mathbf{T}(2k, \Delta)$. This contradicts to the definition of T^* . \square

Lemma 3.9 ([3]). *Let $G(k; \ell)$ be the graph obtained from a non-trivial connected graph G by attaching pendant paths of lengths k and ℓ at some vertex v . If $k \geq \ell \geq 1$, then $\mu_1(Q(G(k; \ell))) > \mu_1(Q(G(k+1; \ell-1)))$.*

Lemma 3.10. *Each path attached at the vertex w of T^* has length at most 2.*

Proof. First we will prove that each path attached at the vertex w of T^* has length at most 3.

Suppose to the contrary that $w_1w_2w_3w_4 \cdots w_s w$ is one path attached at w , where w_1 is a pendant vertex, and $s \geq 4$. Then by Lemma 3.9 it follows that the tree $T' = T^* - w_2w_1 + w_4w_1$ has a larger Laplacian spectral radius. While T' is also in $\mathbf{T}(2k, \Delta)$, and this contradicts to the definition of T^* .

Now we will prove that each path attached at the vertex w of T^* has length at most 2.

Suppose to the contrary that $w w_{13} w_{12} w_{11}$ is a path of length 3 attached at w . Since T^* has perfect matching, $w w_{13} w_{12} w_{11}$ is the unique path with odd length attached at w . And noting the hypothesis $\Delta \geq 3$, we may suppose $w w_{22} w_{21}$ is a path of length 2 attached at w . Then we may apply the following transformations to T^* .

Set

$$T' = \begin{cases} T^* - w_{13}w_{12} + w_{22}w_{12}, & \text{if } x_{w_{22}} \geq x_{w_{13}}; \\ T^* - w_{22}w_{21} + w_{13}w_{21}, & \text{if } x_{w_{13}} > x_{w_{22}}. \end{cases}$$


Then $\mu(T') > \mu(T^*)$ from Lemma 3.4, while T' is also in $\mathbf{T}(2k, \Delta)$. This contradicts to the definition of T^* . So each path attached at the vertex w of T^* has length at most 2. \square

By the similar arguments of Lemma 3.10, we obtain the following results.

Lemma 3.11. *If T^* has a vertex, say u , with degree $2 < d(u) < \Delta$, then each path attached at the vertex u of T^* has length at most 2.*

Combining Lemmas 3.6–3.11, we obtain the main results of this section.

Theorem 3.1. *Let T be a tree in $\mathbf{T}(2k, \Delta)$, and $T^*(2k, \Delta)$ be the tree as shown in Fig. 2. When $\Delta \geq \lceil \frac{k}{2} \rceil + 1$ ($k \geq 4$), we have $\mu(T) \leq \mu(T^*(2k, \Delta))$, and equality holds if and only if $T \cong T^*(2k, \Delta)$.*

Fig. 2. The tree $T^*(2k, \Delta)$.

4. A relation between $\mu(T)$ and $\Delta(T)$ of a tree T in $\mathbf{T}(2k)$

In this section, we will prove that for two trees T_1 and T_2 in $\mathbf{T}(2k)$ ($k \geq 4$), if $\Delta(T_1) > \Delta(T_2)$ and $\Delta(T_1) \geq \lceil \frac{k}{2} \rceil + 2$, then $\mu(T_1) > \mu(T_2)$.

Lemma 4.1 ([1]). Let $G' = G + e$ be the graph obtained from G by inserting a new edge e . Then Laplacian eigenvalues of G interlace the Laplacian eigenvalues of G' , i.e.,

$$\mu_1(G') \geq \mu_1(G) \geq \mu_2(G') \geq \cdots \geq \mu_n(G') \geq \mu_n(G).$$

Lemma 4.2. Let $T^*(2k, \Delta)$ be the tree as shown in Fig. 2. Then $\mu(T^*(2k, \Delta)) < \Delta + 2$, where $\lceil \frac{k}{2} \rceil + 1 \leq \Delta \leq k$, $k \geq 4$.

Proof. First we prove that $\mu_2(T^*(2k, \Delta)) < \Delta + 1$. Denote by $\widehat{K_{1,n-1}}$ the tree (on $2n$ vertices) obtained from the star $K_{1,n-1}$ by attaching an edge at each vertex. Let uv be the edge of $T^*(2k, \Delta)$ (see Fig. 2). Then

$$T^*(2k, \Delta) - uv = \widehat{K_{1,k-\Delta}} \cup \widehat{K_{1,\Delta-2}}.$$

And

$$\mu(T^*(2k, \Delta) - uv) = \max\{\mu(\widehat{K_{1,k-\Delta}}), \mu(\widehat{K_{1,\Delta-2}})\}.$$

It is easy to check that $\max\{d(v) + m(v) \mid v \in V(\widehat{K_{1,k-\Delta}})\} < \Delta + 1$, and $\max\{d(v) + m(v) \mid v \in V(\widehat{K_{1,\Delta-2}})\} < \Delta + 1$. Hence from Lemmas 2.7 and 4.1 we have $\mu_2(T^*(2k, \Delta)) \leq \mu(T^*(2k, \Delta) - uv) < \Delta + 1$.

Let

$$h(x) = x^6 - (k+8)x^5 + (k\Delta - \Delta^2 + 2\Delta + 5k + 23)x^4 - (4k\Delta - 4\Delta^2 + 8\Delta + 9k + 28)x^3 \\ + (5k\Delta - 5\Delta^2 + 10\Delta + 7k + 13)x^2 - (2k\Delta - 2\Delta^2 + 4\Delta + 3k + 2)x + k.$$

Using the similar method of Lemma 2.3, and taking the edge uv of $T^*(2k, \Delta)$ as the edge uv of G in Lemma 2.2, we have

$$\Phi(T^*(2k, \Delta); x) = x(x-2)(x^2-3x+1)^{k-4}h(x).$$

Then $\mu(T^*(2k, \Delta))$ is the largest root of the equation $h(x) = 0$. Noting the fact that $\Delta \geq 3$, we have

$$h(\Delta+1) = -[\Delta^2(\Delta^2 - 2\Delta - 2) + k\Delta + 2\Delta + 1] \\ < 0,$$

and combining the fact that $2\Delta > k+2$, we have

$$h(\Delta+2) = 2\Delta^5 - k\Delta^4 + 6\Delta^4 - 4k\Delta^3 + 4\Delta^3 - 5k\Delta^2 + \Delta^2 - 3k\Delta + 2\Delta - k \\ > 8\Delta^4 - 4k\Delta^3 + 4\Delta^3 - 5k\Delta^2 + \Delta^2 - 3k\Delta + 2\Delta - k \\ > 12\Delta^3 - 5k\Delta^2 + \Delta^2 - 3k\Delta + 2\Delta - k \\ > 12\Delta^2 - 3k\Delta + 2\Delta - k \\ = 3\Delta(4\Delta - k) + (2\Delta - k) \\ > 0.$$

Then there exists a root of the equation $h(x) = 0$ in the interval $(\Delta+1, \Delta+2)$. Note that $\mu_2(T^*(2k, \Delta)) < \Delta+1$, so we know $\mu(T^*(2k, \Delta)) < \Delta+2$. Hence the desired result holds. \square

Lemma 4.3. Let T be a tree in $\mathbf{T}(2k)$ with $\Delta(T) \leq \lceil \frac{k}{2} \rceil$. Then $\mu(T) < \lceil \frac{k}{2} \rceil + 3$.

Proof. Let v be a vertex of T , and write $d(v) = s$ for short. We distinguish the following two cases.

Case 1. $s = 1$.

We have

$$d(v) + m(v) = 1 + m(v) \leq 1 + \left\lceil \frac{k}{2} \right\rceil < \left\lceil \frac{k}{2} \right\rceil + 3. \quad (3.6)$$

Case 2. $2 \leq s \leq \lceil \frac{k}{2} \rceil$.

Let

$$\begin{aligned} N(v) &= \{v_1, v_2, \dots, v_s\}, \\ A(v_i) &= N(v_i) \setminus \{v\}, \quad i = 1, 2, \dots, s. \end{aligned}$$

Then $A(v_1), A(v_2), \dots, A(v_s)$ are pairwise disjoint since T is a tree. And note the fact that T has a perfect matching, then

$$\left| \bigcup_{i=1}^s A(v_i) \right| \leq \frac{2k - (s+1) - (s-1)}{2} + s - 1 = k - 1.$$

So

$$\sum_{i=1}^s |A(v_i)| = \left| \bigcup_{i=1}^s A(v_i) \right| \leq k - 1.$$

Then we have

$$\sum_{u \in N(v)} d(u) \leq s + k - 1.$$

Hence

$$d(v) + m(v) \leq s + \frac{s + k - 1}{s} = s + 1 + \frac{k - 1}{s}.$$

Let $f(s) = s + 1 + \frac{k-1}{s}$. Then $f(s) = s + 1 + \frac{k-1}{s}$ is convex when $s > 0$. Hence when $2 \leq s \leq \lceil \frac{k}{2} \rceil$, we have

$$f(s) \leq \max \left\{ f(2), f\left(\left\lceil \frac{k}{2} \right\rceil\right) \right\} < \left\lceil \frac{k}{2} \right\rceil + 3.$$

So when $2 \leq s \leq \lceil \frac{k}{2} \rceil$, we have

$$d(v) + m(v) < \left\lceil \frac{k}{2} \right\rceil + 3. \quad (3.7)$$

Combining (3.6), (3.7) and Lemma 2.7 we have $\mu(T) < \lceil \frac{k}{2} \rceil + 3$. \square

Now we are ready to obtain our main result of this section.

Theorem 4.1. Let T_1, T_2 be trees in $\mathbf{T}(2k)$ ($k \geq 4$). If $\Delta(T_1) > \Delta(T_2)$ and $\Delta(T_1) \geq \lceil \frac{k}{2} \rceil + 2$, then $\mu(T_1) > \mu(T_2)$.

Proof. We distinguish the following two cases.

Case 1. $\Delta(T_2) \geq \lceil \frac{k}{2} \rceil + 1$.

From Lemma 2.4, 4.2 and Theorem 3.1, we have

$$\mu(T_1) \geq \Delta(T_1) + 1 \geq \Delta(T_2) + 2 > \mu(T^*(2k, \Delta(T_2))) \geq \mu(T_2),$$

where $T^*(2k, \Delta(T_2))$ is the tree defined in Theorem 3.1.

Case 2. $\Delta(T_2) \leq \lceil \frac{k}{2} \rceil$.

From Lemmas 2.4 and 4.3 we have

$$\mu(T_1) \geq \Delta(T_1) + 1 \geq \left\lceil \frac{k}{2} \right\rceil + 3 > \mu(T_2).$$

The proof is completed. \square

Acknowledgements

We would like to thank the referees for her/his careful reading and the valuable suggestions, especially, the proofs of Lemmas 2.8 and 3.10 are simplified according to these suggestions. The first author's research is supported by National Science Foundation of China (No. 10926085), and by Shanghai Leading Academic Discipline Project (No. S30104).

References

- [1] D. Cvetković, P. Rowlinson, S. Simić, *Eigenspaces of Graphs*, Cambridge University Press, 1997.
- [2] D. Cvetković, P. Rowlinson, S. Simić, Signless Laplacians of finite graphs, *Linear Algebra Appl.* 423 (2007) 155–171.
- [3] D. Cvetković, S. Simić, Towards a spectral theory of graphs based on signless Laplacian: I, *Publ. Inst. Math. (N.S.)* 85 (99) (2009) 1–15.
- [4] D. Cvetković, S. Simić, Towards a spectral theory of graphs based on signless Laplacian: II, *Linear Algebra Appl.* 432 (2010) 2257–2272.
- [5] L.H. Feng, Q. Li, X.D. Zhang, Minimizing the Laplacian spectral radius of trees with given matching number, *Linear Multilinear Algebra* 55 (2007) 199–207.
- [6] L.H. Feng, G. Yu, A. Ilić, The Laplacian spectral radius for unicyclic graphs with given independence number, *Linear Algebra Appl.* 433 (2010) 934–944.
- [7] R. Grone, R. Merris, V.S. Sunder, The Laplacian spectrum of a graph, *SIAM J. Matrix Anal. Appl.* 11 (2) (1990) 218–238.
- [8] J.M. Guo, On the Laplacian spectral radius of a tree, *Linear Algebra Appl.* 368 (2003) 379–385.
- [9] J.M. Guo, On the second largest Laplacian eigenvalue of the trees, *Linear Algebra Appl.* 404 (2005) 251–261.
- [10] A.J. Hoffman, J.H. Smith, On the spectral radii of topological equivalent graphs, in: M. Fielder (Ed.), *Recent Advances in Graph Theory*, Academia Praha, Prague, 1975, pp. 273–281.
- [11] Y. Hong, X.D. Zhang, Sharp upper and lower bounds for largest eigenvalue of the Laplacian matrices of trees, *Discrete Math.* 296 (2005) 187–197.
- [12] A. Ilić, On the ordering of trees by the Laplacian coefficients, *Linear Algebra Appl.* 431 (2009) 2203–2212.
- [13] A. Ilić, Trees with minimal Laplacian coefficients, *Comput. Math. Appl.* 59 (2010) 2776–2783.
- [14] R. Merris, A note on Laplacian graph eigenvalue, *Linear Algebra Appl.* 285 (1998) 33–35.
- [15] R. Merris, Laplacian matrices of graphs: a survey, *Linear Algebra Appl.* 197–198 (1994) 143–176.
- [16] J.Y. Shao, L. Zhang, X.Y. Yuan, On the second Laplacian eigenvalues of trees of odd order, *Linear Algebra Appl.* 419 (2006) 475–485.
- [17] X.Y. Yuan, B.F. Wu, E.L. Xiao, The modifications of trees and the Laplacian spectrum, *J. East China Norm. Univ. Natur. Sci. (2)* (2004) 13–18.
- [18] X.Y. Yuan, H.Y. Shan, Y. Liu, On the Laplacian spectral radii of trees, *Discrete Math.* 309 (2009) 4241–4246.
- [19] A.M. Yu, M. Lu, Laplacian spectral radius of trees with given maximum degree, *Linear Algebra Appl.* 429 (2008) 1962–1969.
- [20] X.D. Zhang, The Laplacian spectral radii of trees with degree sequences, *Discrete Math.* 308 (2008) 3143–3150.

Further reading

- [1] R. Grone, R. Merris, Ordering trees by algebraic connectivity, *Graphs Combin.* 6 (1990) 229–237.